

General Information

Nickel MSC is designed for use as a foliar spray on peanuts and other crops where the addition of nickel, copper, and manganese would be beneficial.

Nickel MSC can be tanked mixed with other nutrient and pest control products. Nickel MSC is not recommended for soil application.

Nickel MSC will disperse in water with little agitation. Many pesticides can be added and applied while spraying Nickel MSC. Follow this mixing sequence:

1. Water
2. Nickel MSC
3. Pesticide

Storage Instructions

Do not contaminate water, food or feed by storage or disposal. Store in cool, dry place. May be stored in unheated facility. Nickel MSC has low corrosiveness. Spills from broken containers can be cleaned up by absorbing on clay or other suitable absorbent. Dispose of empty containers according to approved local practices.

Warranty Statement: NIPAN LLC warrants that this product conforms to the chemical description on the label thereof and is reasonably fit for the purpose stated on the label only when used in accordance with directions under normal use conditions. It is impossible to eliminate all risks inherently associated with use of this product. Neither this warranty nor any other warranty of merchantability of fitness of a particular product expressed or implied, extends to the use of this product contrary to label instructions or under abnormal conditions not reasonably foreseeable to the seller, and the buyer assumes the risk of any such use.

Information about the components of this product may be obtained by writing to the manufacturer.



Nickel MSC™

for peanuts and other legumes

Guaranteed Analysis

Sulfur (S)	4.1%
4.1% Combined Sulfur (S)	
Copper (Cu)	1.5%
Manganese (Mn).....	4.5%
Nickel (Ni).....	0.5%

Derived from Copper Sulfate, Manganese Sulfate and Nickel Sulfate and lignosulfonate.

Weight per gallon: 10.1 lbs. 1.21 kg/lr

Net Weight: 25.25 lbs. 11.46 kg

KEEP OUT OF REACH OF CHILDREN

WARNING:



PRECAUTIONARY STATEMENTS

Harmful if swallowed. Avoid contact with skin, eyes, or clothing. Avoid breathing spray mist. May cause irritation of nose, throat and/or skin. Wear goggles and gloves when handling.

FIRST AID

In case of contact, immediately remove clothing and flush skin with plenty of water for several minutes. If irritation persists get medical attention. If product gets into eyes, rinse cautiously with water for several minutes. If swallowed, give large amounts of water. Do not induce vomiting and get medical attention.

Directions For Use

Peanuts:

To increase root mass, peg strength, nodulation, disease resistance to rust, and yield. Improved nickel nutrition of peanuts grown for seed can improve seed germination of saved seed.

Rate and timing: Ground application: Apply 16-24 oz per acre 50 days after planting. Repeat 14 to 21 days later for remainder of the season.

The addition of a non-ionic surfactant at the labeled rate is recommended to improve spray coverage.

Soybeans:

To increase root mass, nodulation and disease resistance to rust. Improved nickel nutrition of soybeans grown for seed can improve seed germination of saved seed.

Rate and timing: Ground application: Apply 12-16 oz in 15-20 gallons water per acre when soybeans are in R1 stage and again 14 to 21 days later.

Other crops:

Nickel MSC can be used in a balanced fertility program where supplemental micronutrient nutrition may improve growth, quality, disease resistance and yield. Improved nickel nutrition has been shown to increase disease resistance to rust.

MANUFACTURED IN THE USA BY: NIPAN L.L.C.

P.O. BOX 5611, 2867 CARROLL ULMER RD.
VALDOSTA, GA 31603-5611.

™ NICKEL MSC IS A TRADEMARK OF NIPAN LLC.

Net Contents: 2.5 Gallons / 9.46 Liters
Case: 2 x 2.5 Gallons / 18.92 Liters

CPC 55723